

# Lift-and-Project Integrality Gaps for the Traveling Salesperson Problem

Thomas Watson\*

Received December 30, 2013; Revised February 28, 2014, final form received March 14, 2014; Published March 28, 2014

**Abstract:** We study the lift-and-project procedures of Lovász-Schrijver and Sherali-Adams applied to the standard linear programming relaxation of the traveling salesperson problem with triangle inequality. For the asymmetric TSP tour problem, Charikar, Goemans, and Karloff (FOCS 2004) proved that the integrality gap of the standard relaxation is at least 2. We prove that after one round of the Lovász-Schrijver or Sherali-Adams procedures, the integrality gap of the asymmetric TSP tour problem is at least  $3/2$ , with a caveat on which version of the standard relaxation is used. For the symmetric TSP tour problem, the integrality gap of the standard relaxation is known to be at least  $4/3$ , and Cheung (SIAM J. Optim. 2005) proved that it remains at least  $4/3$  after  $o(n)$  rounds of the Lovász-Schrijver procedure, where  $n$  is the number of nodes. For the symmetric TSP path problem, the integrality gap of the standard relaxation is known to be at least  $3/2$ , and we prove that it remains at least  $3/2$  after  $o(n)$  rounds of the Lovász-Schrijver procedure, by a simple reduction to Cheung’s result.

## 1 Introduction

The traveling salesperson problem (TSP) is the following: Given a complete directed graph with nonnegative edge distances satisfying the triangle inequality, the goal is to find a shortest route that visits all the nodes. The TSP is one of the most fundamental and well-studied problems in combinatorial optimization

---

\*This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. DGE-0946797 and by the National Science Foundation under Grant No. CCF-1017403.

**Key words and phrases:** lift-and-project, integrality gaps, traveling salesperson problem

(with whole books devoted to it [28, 3, 21]), and there are many variants. For example, in the *tour* version the goal is to find a shortest hamiltonian cycle, while in the *path* version we are additionally given two nodes  $s$  and  $t$  and the goal is to find a shortest hamiltonian path from  $s$  to  $t$ . Along a different dimension, in the *symmetric* version we assume that for every pair of nodes  $u$  and  $v$  the distance from  $u$  to  $v$  equals the distance from  $v$  to  $u$ , while in the *asymmetric* version we make no such assumption.

The approximability of the TSP is not well-understood, and there are large gaps in the approximation ratios between the best NP-hardness results and the best approximation algorithms. The symmetric tour and path versions are shown to be NP-hard to approximate within  $220/219 - \varepsilon$  (for every constant  $\varepsilon > 0$ ) in [38]; this has been improved to  $185/184 - \varepsilon$  in [27], and further to  $123/122 - \varepsilon$  in [25]. The asymmetric tour and path versions are shown to be NP-hard to approximate within  $117/116 - \varepsilon$  (for every constant  $\varepsilon > 0$ ) in [38]; this has been improved to  $75/74 - \varepsilon$  in [25]. The state-of-the-art polynomial-time approximation algorithms achieve approximation ratios  $3/2$  for the symmetric tour version [13],  $8/5$  for the symmetric path version [40] (for a long time the best ratio was  $5/3$  [24], which was later improved to  $\frac{1+\sqrt{5}}{2}$  [2]), and  $O(\log n / \log \log n)$  for the asymmetric tour and path versions [6, 15] where  $n$  is the number of nodes. There are better algorithms for several important restricted classes of distance functions. For example, there have been breakthroughs for shortest path metrics on unweighted undirected graphs: For the symmetric tour version, a  $(3/2 - \varepsilon_0)$ -approximation (for some constant  $\varepsilon_0 > 0$ ) is obtained in [37], further improved to  $1.461$  [31], further improved to  $13/9$  [32], and further improved to  $7/5$  [41], and for the symmetric path version, a  $(5/3 - \varepsilon_0)$ -approximation is obtained in [37, 2], further improved to  $1.586$  [31], further improved to  $19/12 + \varepsilon$  (for every constant  $\varepsilon > 0$ ) [32], further improved to  $1.578$  [2], and further improved to  $3/2$  [41, 18]. The symmetric tour version can be approximated within  $1 + \varepsilon$  (for every constant  $\varepsilon > 0$ ) for Euclidean distances in constant-dimensional real space [4, 30] and for shortest path metrics on weighted undirected planar graphs [5, 26]. The asymmetric tour version has a  $O(1)$ -approximation algorithm for shortest path metrics on weighted directed planar graphs [36].

For problems such as TSP where the known NP-hardness lower bounds on approximation ratios are quite weak, a natural goal is to get stronger lower bounds for restricted classes of algorithms. Such lower bounds can sometimes be construed as evidence that the problem is indeed hard to approximate within the stronger bound, and they may have the additional advantage of being unconditional (NP-hardness lower bounds are conditional on  $P \neq NP$ ). For many combinatorial optimization problems, one very general class of algorithms is those that solve some linear programming relaxation of the problem, “round” the solution to an integral solution in some way, and derive their approximation guarantee by comparing the value of the rounded solution to the optimal value of the relaxation. The integrality gap of an instance is the ratio between the optimal values of the original problem and the relaxation. The existence of a family of instances with integrality gap at least  $\alpha > 1$  proves that no such rounding algorithm can achieve approximation ratio better than  $\alpha$ . Thus exhibiting instances with large integrality gaps constitutes an unconditional lower bound against this class of algorithms. The integrality gap of a relaxation for a problem is defined to be the maximum integrality gap over all instances (expressed as a function of instance size).

For the above variants of the traveling salesperson problem, there is a classic and well-studied linear programming relaxation which has been called the Dantzig-Fulkerson-Johnson relaxation (after [14]), the Held-Karp relaxation (after [22, 23]), and the subtour elimination relaxation. Since there is no consensus on what to call it, we simply refer to it as the *standard relaxation*. Many of the known approximation

algorithms for TSP work by (implicitly or explicitly) rounding solutions to the standard relaxation. For the symmetric tour version, the integrality gap is known to be at least  $4/3$ . It is conjectured to be exactly  $4/3$ , and proving this is a notorious open problem in combinatorial optimization. The best upper bound on the integrality gap is  $3/2$  [45, 43], though in the case of shortest path metrics on unweighted undirected graphs the upper bound has been improved to  $7/5$  [41]. For the symmetric path version, the integrality gap is known to be at least  $3/2$  and at most  $8/5$  [40], though in the case of shortest path metrics on unweighted undirected graphs the upper bound has been improved to  $3/2$  [41, 18]. For the asymmetric tour version, the integrality gap is known to be at least  $2$  [8] and at most  $O(\log n / \log \log n)$  [6]. For the asymmetric path version, the integrality gap is known to be at least  $2$  [8, 16] and at most  $O(\log n / \log \log n)$  [16] (it was earlier shown to be at most  $O(\sqrt{n})$  in [34], and at most  $O(\log n)$  in [17]).

Although integrality gap lower bounds rule out algorithms that derive their approximation guarantees by comparing the output to the optimum of a particular relaxation of the problem, one can always construct tighter and tighter relaxations in hope of reducing the integrality gap and getting improved approximation guarantees. Hence it is desirable to prove lower bounds for large *classes* of relaxations, since this constitutes stronger evidence for the hardness of the problem. Lovász and Schrijver [29] and Sherali and Adams [42] introduced powerful and general techniques for constructing hierarchies of tighter and tighter relaxations for any problem (see [44, 12] for surveys). These techniques are called “lift-and-project” procedures, and there has been substantial interest in proving integrality gap lower bounds for various relaxations produced by these procedures (a small sample of results includes [1, 39, 19, 9, 7]). A nice feature of such results is that lower bounds for sufficiently many rounds of the procedure allow us to rule out slightly-subexponential time rounding algorithms. Not all results on lift-and-project procedures are negative; see, for example, [20] for an application of the Sherali-Adams procedure to obtaining Sparsest Cut algorithms.

We consider the task of proving integrality gap lower bounds for lift-and-project procedures applied to the standard relaxation of the TSP. There is only one previous paper on this topic: Cheung [11] proved an essentially tight result for the Lovász-Schrijver procedure applied to the symmetric TSP tour problem — he showed that the integrality gap remains at least  $4/3$  after  $o(n)$  rounds of the procedure for  $n$ -node instances. A natural question is whether the integrality gap lower bound of  $2$  for the asymmetric TSP tour problem [8] survives these lift-and-project procedures. Our main theorem is a positive result in this direction: We prove that the integrality gap is at least  $3/2$  after one round of the Lovász-Schrijver procedure (which is equivalent to one round of the Sherali-Adams procedure). There is a caveat though: There are two versions of the standard relaxation which are both widely used in the literature and which are equivalent in a certain sense, but one seems to become somewhat weaker than the other after lift-and-project procedures are applied. We can only prove our lower bound for the weaker version (see Section 1.2 for details). Two years after our result was announced, Cheriyan, Gao, Georgiou, and Singla [10] have claimed results that subsume ours and give nontrivial lower bounds for more than one round of the Sherali-Adams procedure, though their results give nothing new for the Lovász-Schrijver procedure. We discuss this follow-up work in Section 1.2.

One key challenge in proving Lovász-Schrijver integrality gap lower bounds is in designing so-called protection matrices. None of the previous techniques for designing protection matrices (for problems such as vertex cover, constraint satisfaction problems, and symmetric TSP) seem to help with asymmetric TSP. We introduce a new technique for our setting, based on finding certain combinatorial objects we call

*frames* in directed graphs.

Another natural question is whether the folklore integrality gap lower bound of  $3/2$  for the symmetric TSP path problem survives lift-and-project procedures. We show that it does, in fact, survive  $o(n)$  rounds of the Lovász-Schrijver procedure, by giving a simple reduction to Cheung's result [11]. The reduction is not generic, but rather exploits specific properties of Cheung's family of instances. This result can be considered evidence that the symmetric TSP path problem cannot be approximated better than  $3/2$  (even by slightly-subexponential time algorithms). The fact that we get a lower bound of  $3/2$  in both this result and our result for the asymmetric TSP tour problem is a coincidence; the techniques used are quite different.

## 1.1 Definitions

In Section 1.1.1 we give the definitions of integrality gaps and the Lovász-Schrijver lift-and-project procedure. In Section 1.1.2 we give the definitions of the relevant variants of the traveling salesperson problem and their standard linear programming relaxations.

### 1.1.1 Lift-and-Project Integrality Gaps

For our purpose a (*combinatorial optimization*) *problem* is a set of instances, where each instance has a set of points  $I \subseteq \{0, 1\}^m$  and a nonnegative convex objective function  $f$ , and the goal is to minimize  $f$  over points in  $I$ . In the notation, we suppress the dependence of  $I$ ,  $m$ , and  $f$  on the instance. The instances of a problem may have some size parameter  $n$  (not necessarily equal to  $m$ ). A *relaxation* of a problem associates to each instance a convex set  $R \subseteq [0, 1]^m$  such that  $I \subseteq R$ . Given a relaxation of a problem, the *integrality gap* of an instance is defined to be  $\frac{\min_{x \in I} f(x)}{\min_{x \in R} f(x)}$ , and the integrality gap of the relaxation is defined to be the maximum integrality gap of an instance, expressed as a function of the instance size  $n$ . If the objective function for a problem is always linear, and the set  $R$  for a relaxation is always a polytope, then we call the relaxation a *linear programming relaxation*.

Lovász and Schrijver [29] introduced a general technique for improving linear programming relaxations. Given a convex set  $R \subseteq [0, 1]^m$ , the Lovász-Schrijver procedure produces a new convex set  $N(R) \subseteq [0, 1]^m$  which is a polytope if  $R$  is. The procedure can be applied iteratively, and we use  $N^r(R)$  to denote the convex set after  $r$  rounds (for example  $N^2(R) = N(N(R))$ ). The  $N$  operator has the following properties:  $R \supseteq N(R) \supseteq \text{conv}(R \cap \{0, 1\}^m)$  and  $N^m(R) = \text{conv}(R \cap \{0, 1\}^m)$  where  $\text{conv}(\cdot)$  denotes convex hull, and thus

$$R \supseteq N(R) \supseteq N^2(R) \supseteq \cdots \supseteq N^{m-1}(R) \supseteq N^m(R) = \text{conv}(R \cap \{0, 1\}^m).$$

Hence the procedure can be applied to a relaxation of a combinatorial optimization problem to yield new relaxations, where the integrality gap is nonincreasing with the number of rounds and drops to 1 after at most  $m$  rounds if  $R \cap \{0, 1\}^m = I$ . Furthermore, if  $R$  has a polynomial-time separation oracle, then linear objectives can be optimized over  $N^r(R)$  in time  $m^{O(r)}$ .

We now define the  $N$  operator. Given a convex set  $R \subseteq [0, 1]^m$  we define

$$\text{cone}(R) = \{(\lambda, \lambda x_1, \dots, \lambda x_m) : \lambda \geq 0 \text{ and } (x_1, \dots, x_m) \in R\}$$

and we index the  $\lambda$  coordinate by 0. For an  $(m+1) \times (m+1)$  matrix  $X$  we index the top row and left column by 0, and we use  $X_i$  to denote the  $i$ th row ( $i \in \{0, 1, \dots, m\}$ ). Given a convex set  $R \subseteq [0, 1]^m$ , a point  $x \in [0, 1]^m$  is in  $N(R)$  if and only if there exists an  $(m+1) \times (m+1)$  matrix  $X$ , called a *protection matrix*, such that

- (i)  $X$  is symmetric,
- (ii)  $X_0 = \text{diag}(X) = (1 \ x)$ , and
- (iii)  $X_i \in \text{cone}(R)$  and  $X_0 - X_i \in \text{cone}(R)$  for each  $i \in \{1, \dots, m\}$ .

The procedure is called a “lift-and-project” procedure because it first lifts  $R$  to a convex set in  $[0, 1]^{(m+1)^2}$  (namely the set of matrices satisfying (i), (ii), and (iii) for some  $x$ ) which is a polytope if  $R$  is, and then projects on certain coordinates to get  $N(R)$ . Lovász and Schrijver also introduced a stronger operator  $N_+$  which is defined in the same way as  $N$  except the protection matrix is also required to be positive semidefinite. For all convex sets  $R \subseteq [0, 1]^m$  we have  $N_+(R) \subseteq N(R)$  and

$$R \supseteq N_+(R) \supseteq \dots \supseteq N_+^m(R) = \text{conv}(R \cap \{0, 1\}^m).$$

Sherali and Adams [42] introduced a lift-and-project procedure for improving linear programming relaxations which is more powerful than the Lovász-Schrijver  $N$  procedure, but also more difficult to prove integrality gap lower bounds for. One round of Sherali-Adams coincides with one round of Lovász-Schrijver. We do not define the procedure for higher rounds here.

### 1.1.2 The Traveling Salesperson Problem

We now formally define the traveling salesperson problem and its standard linear programming relaxation [14, 22, 23]. For completeness, we define all four variants discussed above. Given an undirected graph  $G = (V, E)$  and a set  $S \subseteq V$  we let  $\delta(S)$  denote the set of edges crossing the cut  $(S, \bar{S})$ . For  $v \in V$  we define  $\delta(v)$  to be  $\delta(\{v\})$ . Given a directed graph  $G = (V, E)$  and a set  $S \subseteq V$  we let  $\delta^+(S)$  denote the set of edges leaving  $S$  and  $\delta^-(S)$  denote the set of edges entering  $S$ . For  $v \in V$  we define  $\delta^+(v)$  to be  $\delta^+(\{v\})$  and  $\delta^-(v)$  to be  $\delta^-(\{v\})$ . For every graph  $G = (V, E)$  (undirected or directed), given a set  $F \subseteq E$  and a vector  $x = (x_e)_{e \in E}$  we define  $x(F) = \sum_{e \in F} x_e$ . Also, given two vectors  $d = (d_e)_{e \in E}$  and  $x = (x_e)_{e \in E}$  we define  $d \cdot x = \sum_{e \in E} d_e x_e$ . We say a directed graph is complete if every pair of nodes has both edges between them and there are no self loops. We use  $K_n$  to denote both the complete undirected graph on  $n$  nodes and the complete directed graph on  $n$  nodes; it will always be clear from context which is meant. We always use  $n$  for the number of nodes in a graph, and  $m$  for the number of variables (as in Section 1.1.1) which, for our purpose, is always the number of edges.

**Symmetric tour version.** Given an undirected graph  $G = (V, E)$ , the *symmetric tour polytope*  $ST(G)$  has variables  $x = (x_e)_{e \in E}$  and is defined by the following constraints.

$$\begin{aligned} x(\delta(S)) &\geq 2 & \forall \emptyset \subsetneq S \subsetneq V \\ x(\delta(v)) &= 2 & \forall v \in V \\ x_e &\in [0, 1] & \forall e \in E \end{aligned}$$

We define  $ST^{\text{int}}(G)$  similarly but require  $x_e \in \{0, 1\}$  for all  $e \in E$ ; note that  $ST^{\text{int}}(G)$  consists of exactly the hamiltonian cycles in  $G$ . The *symmetric TSP tour problem* is the following: Given a complete undirected graph  $K_n = (V, E)$  with nonnegative edge distances  $d = (d_e)_{e \in E}$  satisfying the triangle inequality, minimize  $d \cdot x$  subject to  $x \in ST^{\text{int}}(K_n)$ . The standard relaxation allows  $x \in ST(K_n)$ .

**Symmetric path version.** Given an undirected graph  $G = (V, E)$  and distinct nodes  $s, t \in V$  (the case  $s = t$  is covered by the symmetric tour version), the *symmetric path polytope*  $SP(G, s, t)$  has variables  $x = (x_e)_{e \in E}$  and is defined by the following constraints.

$$\begin{aligned} x(\delta(S)) &\geq 2 & \forall \emptyset \subsetneq S \subsetneq V \text{ with } |S \cap \{s, t\}| \neq 1 \\ x(\delta(S)) &\geq 1 & \forall \emptyset \subsetneq S \subsetneq V \text{ with } |S \cap \{s, t\}| = 1 \\ x(\delta(v)) &= 2 & \forall v \in V \setminus \{s, t\} \\ x(\delta(v)) &= 1 & \forall v \in \{s, t\} \\ x_e &\in [0, 1] & \forall e \in E \end{aligned}$$

We define  $SP^{\text{int}}(G, s, t)$  similarly but require  $x_e \in \{0, 1\}$  for all  $e \in E$ ; note that  $SP^{\text{int}}(G, s, t)$  consists of exactly the hamiltonian paths in  $G$  with  $s$  and  $t$  as their endpoints. The *symmetric TSP path problem* is the following: Given a complete undirected graph  $K_n = (V, E)$  with nonnegative edge distances  $d = (d_e)_{e \in E}$  satisfying the triangle inequality, and given distinct nodes  $s, t \in V$ , minimize  $d \cdot x$  subject to  $x \in SP^{\text{int}}(K_n, s, t)$ . The standard relaxation allows  $x \in SP(K_n, s, t)$ .

**Asymmetric tour version.** Given a directed graph  $G = (V, E)$ , the *asymmetric tour polytope*  $AT(G)$  has variables  $x = (x_e)_{e \in E}$  and is defined by the following constraints.

$$\begin{aligned} x(\delta^+(S)) &\geq 1 & \forall \emptyset \subsetneq S \subsetneq V \\ x(\delta^-(S)) &\geq 1 & \forall \emptyset \subsetneq S \subsetneq V \\ x(\delta^+(v)) &= 1 & \forall v \in V \\ x(\delta^-(v)) &= 1 & \forall v \in V \\ x_e &\in [0, 1] & \forall e \in E \end{aligned}$$

Of course, the second group of constraints is redundant given the first, but we prefer to include it for aesthetic reasons. We define  $AT^{\text{int}}(G)$  similarly but require  $x_e \in \{0, 1\}$  for all  $e \in E$ ; note that  $AT^{\text{int}}(G)$  consists of exactly the hamiltonian cycles in  $G$ . The *asymmetric TSP tour problem* is the following: Given a complete directed graph  $K_n = (V, E)$  with nonnegative edge distances  $d = (d_e)_{e \in E}$  satisfying the triangle inequality, minimize  $d \cdot x$  subject to  $x \in AT^{\text{int}}(K_n)$ . The standard relaxation allows  $x \in AT(K_n)$ .

**Asymmetric path version.** Given a directed graph  $G = (V, E)$  and distinct nodes  $s, t \in V$  (the case  $s = t$  is covered by the asymmetric tour version), the *asymmetric path polytope*  $AP(G, s, t)$  has variables

$x = (x_e)_{e \in E}$  and is defined by the following constraints.

$$\begin{aligned}
 x(\delta^+(S)) &\geq 1 & \forall \emptyset \subsetneq S \subseteq V \setminus \{t\} \\
 x(\delta^-(S)) &\geq 1 & \forall \emptyset \subsetneq S \subseteq V \setminus \{s\} \\
 x(\delta^+(v)) &= 1 & \forall v \in V \setminus \{t\} \\
 x(\delta^-(v)) &= 1 & \forall v \in V \setminus \{s\} \\
 x(\delta^+(t)) &= 0 \\
 x(\delta^-(s)) &= 0 \\
 x_e &\in [0, 1] & \forall e \in E
 \end{aligned}$$

The importance of the constraints  $x(\delta^+(t)) = x(\delta^-(s)) = 0$  was clarified in [33]. We define  $AP^{\text{int}}(G, s, t)$  similarly but require  $x_e \in \{0, 1\}$  for all  $e \in E$ ; note that  $AP^{\text{int}}(G, s, t)$  consists of exactly the hamiltonian paths in  $G$  from  $s$  to  $t$ . The *asymmetric TSP path problem* is the following: Given a complete directed graph  $K_n = (V, E)$  with nonnegative edge distances  $d = (d_e)_{e \in E}$  satisfying the triangle inequality, and given distinct nodes  $s, t \in V$ , minimize  $d \cdot x$  subject to  $x \in AP^{\text{int}}(K_n, s, t)$ . The standard relaxation allows  $x \in AP(K_n, s, t)$ .

Although the cut constraints in the four polytopes  $ST$ ,  $SP$ ,  $AT$ ,  $AP$  look rather different from each other, they are actually very uniform: Every cut should have capacity at least 2 (symmetric versions) or at least 1 in both directions (asymmetric versions), unless the cut separates  $s$  and  $t$  in the path versions, in which case it should have capacity at least 1 (symmetric version) or at least 1 from  $s$  to  $t$  (asymmetric version). Note that although all four polytopes have exponentially many constraints, they each have a polynomial-time separation oracle using min-cut computations. Hence linear objectives can be optimized over these polytopes in polynomial time using the ellipsoid algorithm.

## 1.2 Results

There is only one previous result on lift-and-project integrality gaps for the traveling salesperson problem: Cheung [11] proved that the integrality gap of the standard relaxation of the symmetric TSP tour problem remains at least  $4/3 - o(1)$  even after  $o(n)$  rounds of the Lovász-Schrijver  $N_+$  procedure.<sup>1</sup> We consider whether a lower bound better than  $4/3$  can be obtained for the *asymmetric* TSP tour problem. Although we cannot prove a better lower bound for the standard relaxation, we can prove a new result for the following relaxation, which is equivalent to the standard relaxation in a certain sense.

Given a directed graph  $G = (V, E)$ , the *balanced asymmetric tour polytope*  $AT_{\text{bal}}(G)$  has variables  $x = (x_e)_{e \in E}$  and is defined by the following constraints.

$$\begin{aligned}
 x(\delta^+(S)) &\geq 1 & \forall \emptyset \subsetneq S \subsetneq V \\
 x(\delta^-(S)) &\geq 1 & \forall \emptyset \subsetneq S \subsetneq V \\
 x(\delta^+(v)) &= x(\delta^-(v)) & \forall v \in V \\
 x_e &\in [0, 1] & \forall e \in E
 \end{aligned}$$

---

<sup>1</sup>Note that the number of variables in the relaxation is  $m = \binom{n}{2}$ , so this result can be phrased as handling  $o(\sqrt{m})$  rounds.

As before, the second group of constraints is redundant given the first, but we prefer to include it for aesthetic reasons. We define  $AT_{\text{bal}}^{\text{int}}(G)$  similarly but require  $x_e \in \{0, 1\}$  for all  $e \in E$ .

We have  $AT(G) \subseteq AT_{\text{bal}}(G)$  and thus the balanced asymmetric tour polytope yields a relaxation for the asymmetric TSP tour problem, which we call the *balanced standard relaxation*. It is well-known that  $AT_{\text{bal}}(K_n)$  is equivalent to  $AT(K_n)$ , and  $AT_{\text{bal}}^{\text{int}}(K_n)$  is equivalent to  $AT^{\text{int}}(K_n)$ , in the sense that they have the same minimum values under any objective  $d \cdot x$  where  $d$  satisfies the triangle inequality (see for example [8, 35]). For this reason, both formulations are very commonly used in the literature and are considered interchangeable. However, they are not necessarily interchangeable after lift-and-project procedures are applied;  $N(AT(K_n))$  might have a smaller integrality gap than  $N(AT_{\text{bal}}(K_n))$ . The integrality gap of  $AT_{\text{bal}}(K_n)$  does drop to 1 after at most  $m = n(n - 1)$  rounds of  $N$ , since optimizing over  $AT_{\text{bal}}^{\text{int}}(K_n)$  is equivalent to optimizing over  $AT^{\text{int}}(K_n)$  in the sense described above. Our lower bound only applies to the balanced standard relaxation. It is not clear how large of a caveat this is, since sometimes insignificant modifications to a linear programming relaxation can turn out to be very significant after lift-and-project procedures are applied [12].

**Theorem 1.1.** *The integrality gap of one round of the Lovász-Schrijver  $N$  procedure applied to the balanced standard relaxation of the asymmetric TSP tour problem is at least  $3/2 - o(1)$ .*

We prove Theorem 1.1 in Section 2. The instances we use to witness Theorem 1.1 are the same as the ones constructed in [8], but we need a new analysis showing that a certain fractional solution survives one round of lift-and-project. The heart of the analysis involves finding certain sets of edges, which we call *frames*, in the graphs constructed in [8]. We need to find a frame associated with each edge, with the property that for every pair of edges  $e_1$  and  $e_2$ ,  $e_2$  is in  $e_1$ 's frame if and only if  $e_1$  is in  $e_2$ 's frame (this property is related to the symmetry requirement for protection matrices).

Note that the lower bound of  $3/2 - o(1)$  beats the lower bound of  $4/3 - o(1)$  that follows trivially from Cheung's result for the symmetric TSP tour problem. There are four deficiencies in Theorem 1.1 that would be nice to overcome.

- (1) We only handle a single round of lift-and-project; ideally we would like to handle at least a superconstant number of rounds.
- (2) We only handle the balanced standard relaxation; ideally we would like to handle the standard relaxation as defined in Section 1.1.2.
- (3) The integrality gap lower bound is only  $3/2 - o(1)$ ; ideally we would like to match the lower bound of  $2 - o(1)$  due to [8] which holds for zero rounds of lift-and-project.
- (4) The lower bound only holds for the Lovász-Schrijver  $N$  procedure; ideally we would like to handle the stronger  $N_+$  procedure.

Although these deficiencies make the result quantitatively somewhat weak, strong lower bounds for the approximability of TSP are quite scarce because the global structure of the problem makes its approximability hard to understand. Our result is one of very few results on lift-and-project procedures applied to problems with global constraints. There seems to be a natural barrier for the basic technique used to prove Theorem 1.1: While the lower bound is  $2/1$  for 0 rounds and  $3/2$  for 1 round, it seems

generalizing the technique would lead to a lower bound of at most  $4/3$  for 2 rounds. However,  $4/3$  is already known by Cheung's result [11], even for  $o(n)$  rounds and even for the symmetric version. Getting a lower bound of  $4/3$  for 2 rounds of the *Sherali-Adams* procedure would be a new result, and indeed in follow-up work [10], Cheriyan, Gao, Georgiou, and Singla accomplished this: They prove that for every  $t \geq 0$ , the integrality gap of  $t$  rounds of the Sherali-Adams procedure applied to the balanced standard relaxation of the asymmetric TSP tour problem is at least  $1 + 1/(t+1) - o(1)$  (also using the instances of [8]), which subsumes Theorem 1.1. They also prove that for every  $t \geq 0$ , the integrality gap of  $t$  rounds of the Sherali-Adams procedure applied to the standard relaxation (not the balanced one) of the asymmetric TSP tour problem is at least  $1 + 1/(2t+3) - o(1)$  (using a new instance construction).

We now turn to the symmetric TSP path problem. By giving a simple reduction to Cheung's result [11], we show that the folklore  $3/2 - o(1)$  lower bound on the integrality gap of the standard relaxation survives  $o(n)$  rounds of the Lovász-Schrijver  $N_+$  procedure. The reduction is not a generic reduction, but rather exploits special properties of Cheung's family of instances.

**Theorem 1.2.** *The integrality gap of  $o(n)$  rounds of the Lovász-Schrijver  $N_+$  procedure applied to the standard relaxation of the symmetric TSP path problem is at least  $3/2 - o(1)$ . Moreover, this lower bound holds even for instances that are shortest path metrics on unweighted undirected graphs.*

We prove Theorem 1.2 in Section 3.

## 2 Asymmetric TSP Tour Problem

In this section we prove Theorem 1.1.

### 2.1 The Instances


For every  $\varepsilon > 0$  we need to construct an instance  $d = (d_e)_{e \in E}$  on  $K_n = (V, E)$  such that

$$\frac{\min_{x \in AT^{\text{int}}(K_n)} d \cdot x}{\min_{x \in N(AT_{\text{bal}}(K_n))} d \cdot x} \geq 3/2 - \varepsilon. \quad (2.1)$$

The instances we use are the same ones constructed by Charikar, Goemans, and Karloff [8]. For integers  $k \geq 1$  and  $r \geq 2$  they construct a directed graph  $G_{k,r}$  with nonnegative edge weights as follows (see the illustrations in Figures 1, 2, and 3). The graph  $G_{1,r}$  consists of two paths of  $r+1$  edges on the same  $r+2$  nodes, going in opposite directions, and all edges have weight 1. One of the two endpoints is designated as the source  $s$  and the other as the sink  $t$ . For  $k > 1$  the graph  $G_{k,r}$  consists of  $r$  copies of  $G_{k-1,r}$  and an additional source node  $s$  and sink node  $t$ , with a path from  $s$  to  $t$  of  $r+1$  edges visiting the sources of the copies in some order, and another path from  $t$  to  $s$  of  $r+1$  edges visiting the sinks of the copies in the opposite order. All the new edges have weight  $r^{k-1}$ .

Next, as in [8] we define a directed graph  $L_{k,r} = (V_{k,r}, E_{k,r})$  with nonnegative edge weights as follows. Suppose  $(s, v_1), (v_2, t), (t, v_3), (v_4, s)$  are the edges incident to  $s$  and  $t$  in  $G_{k,r}$ . Then  $L_{k,r}$  is defined by taking  $G_{k,r}$ , removing  $s$  and  $t$ , and including edges  $(v_2, v_1)$  and  $(v_4, v_3)$ , both of weight  $r^{k-1}$ .

Now fix some  $k \geq 1$  and  $r \geq 2$  and let  $n = |V_{k,r}| = \Theta(r^k)$ . We define the edge distances  $d = (d_e)_{e \in E}$  on the complete directed graph  $K_n = (V_{k,r}, E)$  as the shortest path distances in the weighted graph  $L_{k,r}$ . Note that  $d$  is nonnegative and satisfies the triangle inequality, so this is a valid instance.

Figure 1: The graph  $G_{1,3}$ .Figure 2: The graph  $G_{2,3}$ .

**Lemma 2.1** ([8]). *For  $k \geq 2$  and  $r \geq 3$  we have  $\min_{x \in AT^{int}(K_n)} d \cdot x \geq (2k-1)(r-1)r^{k-1}$ .*

**Lemma 2.2.** *For  $k \geq 1$  and  $r \geq 2$  we have  $\min_{x \in N(AT_{bal}(K_n))} d \cdot x \leq \frac{4}{3}k(r+1)r^{k-1}$ .*

Letting  $k \geq 2$  and  $r \geq 3$  and combining Lemma 2.1 and Lemma 2.2, we find that the ratio on the left side of (2.1) is at least  $\frac{3}{2} \cdot \frac{(k-1/2)(r-1)}{k(r+1)}$ , which is at least  $3/2 - \varepsilon$  provided  $k$  and  $r$  are large enough. This finishes the proof of Theorem 1.1, assuming Lemma 2.2.


## 2.2 The Fractional Solution

We now prove Lemma 2.2. Consider the vector  $x = (x_e)_{e \in E_{k,r}}$  with  $x_e = 2/3$  for all  $e \in E_{k,r}$ . We claim that  $x \in N(AT_{bal}(L_{k,r}))$ . Then a simple argument shows that extending  $x$  with 0 values for all the edges in  $E \setminus E_{k,r}$  yields a point  $\hat{x} \in N(AT_{bal}(K_n))$  such that  $d \cdot \hat{x}$  equals  $2/3$  times the sum of all the edge weights in  $L_{k,r}$  (using the fact that every edge in the weighted graph  $L_{k,r}$  is a shortest path from its tail to its head). An inductive argument shows that the sum of all the edge weights in  $L_{k,r}$  is at most  $2k(r+1)r^{k-1}$  (see [8]), so  $\hat{x}$  witnesses Lemma 2.2.

All that remains is to prove that  $x \in N(AT_{bal}(L_{k,r}))$ , and this is the meat of the argument. In order to exhibit a protection matrix for  $x$ , we first need the following definitions. We call a path or cycle in a directed graph *edge-simple* if it does not repeat any edges, but may repeat nodes.

**Definition 2.3.** Given a directed graph  $G' = (V', E')$  and an edge  $(u, v) \in E'$ , a *frame* for  $(u, v)$  is a set of edges  $F \subseteq E' \setminus \{(u, v)\}$  that consists of an edge-simple path from  $u$  to  $v$  together with zero or more edge-simple cycles. The path and cycles are required to be edge-disjoint from each other.

**Claim 2.4.** *In the graph  $L_{k,r}$  there exists a frame  $F_e$  for each  $e \in E_{k,r}$ , with the property that for all  $e_1, e_2 \in E_{k,r}$  we have  $e_2 \in F_{e_1}$  if and only if  $e_1 \in F_{e_2}$ .*


 Figure 3: The graph  $G_{3,3}$ .

We prove Claim 2.4 shortly, but let us now see how to use it to construct a protection matrix  $X$  for  $x$ . The rows and columns of  $X$  are indexed by  $E_{k,r}$ , except there is an additional 0th row and 0th column. We must have  $X_{0,0} = 1$  and  $X_{0,e} = X_{e,0} = X_{e,e} = 2/3$  for all  $e \in E_{k,r}$ . For  $e_1 \neq e_2$  we set

$$X_{e_1,e_2} = \begin{cases} 1/3 & \text{if } e_2 \in F_{e_1} \\ 1/2 & \text{otherwise} \end{cases}.$$

Note that  $X$  is symmetric by the property in Claim 2.4. So we just need to show that  $X_e \in \text{cone}(AT_{\text{bal}}(L_{k,r}))$  and  $X_0 - X_e \in \text{cone}(AT_{\text{bal}}(L_{k,r}))$  for all  $e \in E_{k,r}$ .

We first consider  $X_e$ . Since  $(X_e)_0 = 2/3$ ,  $X_e \in \text{cone}(AT_{\text{bal}}(L_{k,r}))$  is equivalent to  $x^{(e)} \in AT_{\text{bal}}(L_{k,r})$  where  $x^{(e)}$  is the vector on  $E_{k,r}$  that gives value 1 to  $e$ , value 1/2 to the edges in  $F_e$ , and value 3/4 to the remaining edges. To verify the balance condition  $x^{(e)}(\delta^+(v)) = x^{(e)}(\delta^-(v))$  for all  $v \in V_{k,r}$ , consider starting with the vector that assigns 3/4 to all edges. Since every node in  $L_{k,r}$  has two incoming edges and two outgoing edges, this vector would satisfy the balance condition. Reducing the values on the path and cycles to 1/2 maintains the balance condition except at the endpoints of  $e$ , which are remedied by raising the value of  $e$  to 1. To verify the cut constraints, suppose we interpret  $x^{(e)}$  as capacities in a flow network. Then we need to show that every cut has capacity at least 1 in both directions, which is equivalent to showing that a unit of flow can be sent from any node to any other node. It is shown in [8] that for every  $u, v \in V_{k,r}$  there exist two edge-disjoint paths from  $u$  to  $v$  in  $L_{k,r}$ . Since every edge has capacity at least 1/2 in  $x^{(e)}$ , we can send a half-unit of flow along each of these two paths. Thus we have shown that  $X_e \in \text{cone}(AT_{\text{bal}}(L_{k,r}))$ .

Now we consider  $X_0 - X_e$ . Since  $(X_0 - X_e)_0 = 1/3$ ,  $X_0 - X_e \in \text{cone}(AT_{\text{bal}}(L_{k,r}))$  is equivalent to  $y^{(e)} \in AT_{\text{bal}}(L_{k,r})$  where  $y^{(e)}$  is the vector on  $E_{k,r}$  that gives value 0 to  $e$ , value 1 to the edges in  $F_e$ , and value 1/2 to the remaining edges. The balance condition is verified similarly as before: The vector that assigns 1/2 to all edges would satisfy the balance condition, and raising the values on the path and cycles to 1 maintains the balance condition except at the endpoints of  $e$ , which are remedied by reducing the value of  $e$  to 0. To verify the cut constraints, we need to show that a unit of flow can be sent from any

node to any other node, respecting the capacities  $y^{(e)}$ . Every edge has capacity at least  $1/2$  except for  $e$  itself, so if we try to send a half-unit of flow along each of the two edge-disjoint paths from  $u$  to  $v$ , we only run into trouble if one of these paths uses  $e$ . In that case, the half-unit of flow through  $e$  can be re-routed via the path in the frame  $F_e$ . This path may overlap with the two paths from  $u$  to  $v$ , but since each edge in  $F_e$  has capacity 1, it can accommodate two half-units of flow simultaneously. Thus we can always send a unit of flow from  $u$  to  $v$ , and we have shown that  $X_0 - X_e \in \text{cone}(AT_{\text{bal}}(L_{k,r}))$ .

This finishes the proof of Lemma 2.2, assuming Claim 2.4.

### 2.3 The Frames

We now construct the frames used to prove Claim 2.4, and in the next subsection we verify that these frames satisfy the symmetry property.


For each edge  $e \in E_{k,r}$ , among all paths from  $e$ 's tail to  $e$ 's head that do not use  $e$ , there is a unique shortest such path (having the fewest edges). The natural first attempt at defining the frames is to take  $F_e$  to be this path for each  $e$ . However, this does not work in general; it is crucial that we augment the path with some cycles.


Each edge  $e \in E_{k,r}$  can be assigned a level  $\ell \in \{1, \dots, k\}$ , meaning that  $e$  is introduced as part of a copy of  $G_{\ell,r}$  in the construction of  $G_{k,r}$  (and the two new edges in  $L_{k,r}$  are at level  $k$ ). The most important distinction is whether  $\ell < k$  or  $\ell = k$ . If  $\ell < k$  then we include in  $F_e$  the shortest path from  $e$ 's tail to  $e$ 's head that does not use  $e$ , and for every copy of  $G_{\ell,r}$  that is contained in the ambient copy of  $G_{\ell+1,r}$  (or in  $L_{k,r}$  if  $\ell = k-1$ ) and that is not touched by this path, we include in  $F_e$  a cycle containing all the edges at level  $\ell$  in this copy of  $G_{\ell,r}$ . If  $\ell = k$  then we only include in  $F_e$  the shortest path from  $e$ 's tail to  $e$ 's head that does not use  $e$ , and we include no cycles. This completes the description of the frames.

Let us be a bit more precise and give some illustrations. From now on, we use “terminal” to mean either source or sink, since there is no need to distinguish between the latter. We say an edge at level  $\ell < k$  is “mediating” if it is incident to a terminal when it is introduced in  $G_{\ell,r}$ , and “nonmediating” otherwise. Also, we say an edge at level  $\ell < k-1$  is “outer” if when we start at its head and follow the direct route to a terminal of the ambient copy of  $G_{\ell,r}$ , the other edge entering that terminal is a mediating edge. The edge is an “inner” edge otherwise, and all edges at level  $k-1$  are considered inner.

Consider an edge  $e = (u, v)$  at level  $\ell < k$ . There are four cases depending on whether  $e$  is mediating/nonmediating and outer/inner. The “typical” case is when  $e$  is nonmediating and inner and at level  $3 \leq \ell \leq k-2$ . This is illustrated in Figure 4, with bold frame edges. Then  $u$  is a terminal of a copy of  $G_{\ell-1,r}$ , and the path starts by taking the direct route to the other terminal of this copy, then the direct route to a terminal of the ambient copy of  $G_{\ell,r}$ , then one step to the next copy of  $G_{\ell,r}$ , then the direct route to the other terminal of that copy, then one step back to the copy of  $G_{\ell,r}$  containing  $e$ , then the direct route to the other terminal of the copy of  $G_{\ell-1,r}$  of which  $v$  is a terminal, and finally the direct route to  $v$  through this copy of  $G_{\ell-1,r}$ . This path involves two copies of  $G_{\ell,r}$ . For each of the other  $r-2$  copies of  $G_{\ell,r}$  in the ambient copy of  $G_{\ell+1,r}$ , we include a cycle containing all the edges at level  $\ell$  in this copy of  $G_{\ell,r}$ . This completes the description of the frame  $F_e$ . Nothing changes if  $\ell = 2$  or  $\ell = k-1$ . When  $\ell = 1$ , there are no copies of “ $G_{\ell-1,r}$ ” to go through, so the path simply starts by going directly to a terminal of the copy of  $G_{1,r}$  containing  $e$ , and ends by going from the other terminal directly to  $v$ .

When  $e$  is mediating, all that changes is that one of the two copies of  $G_{\ell-1,r}$  disappears; either  $u$  or  $v$  is already a terminal of the ambient copy of  $G_{\ell,r}$ . When  $e$  is outer, there is no “next” copy of  $G_{\ell,r}$ ; the

Figure 4: Frame for a nonmediating inner edge at level  $3 \leq \ell \leq k - 2$ .

Figure 5: Frame for a mediating outer edge at level  $\ell = 2$ .


first step along an edge at level  $\ell + 1$  takes us to a terminal of the ambient copy of  $G_{\ell+1,r}$ , then the next step takes us back to the copy of  $G_{\ell,r}$  containing  $e$ . Furthermore when  $e$  is outer,  $r - 1$  cycles need to be included in the frame (rather than  $r - 2$ ) since the path only touches one copy of  $G_{\ell,r}$ . Figure 5 illustrates the frame for a mediating outer edge at level  $\ell = 2$ .

Figure 6 illustrates the frame for an edge  $e = (u, v)$  at level  $\ell = k$ . The path takes the direct route from  $u$  to the other terminal of the copy of  $G_{k-1,r}$  containing  $u$ , then follows the level- $k$  cycle containing this terminal until it gets to the copy of  $G_{k-1,r}$  containing  $v$ , then follows the direct route to  $v$  through this copy. This completes the more precise description of all the frames.

## 2.4 Verifying the Symmetry Property

It remains to be verified that for all  $e_1, e_2 \in E_{k,r}$  we have  $e_2 \in F_{e_1}$  if and only if  $e_1 \in F_{e_2}$ . Of course, we just need to consider an arbitrary  $e_1$  and show that for each  $e_2 \in F_{e_1}$  we have  $e_1 \in F_{e_2}$ . This is a slightly tedious case analysis. We first give a couple illustrative cases, which the reader may find to be sufficiently convincing. Suppose  $e_1$  is the edge  $e$  in Figure 4. If  $e_2$  is the third edge on the path in  $F_{e_1}$ , then the path in  $F_{e_2}$  first goes right to the other terminal in the copy of  $G_{\ell-2,r}$ , then goes down to  $e_1$ 's tail, then traverses  $e_1$ , then goes up through the copy of  $G_{\ell-1,r}$  containing  $e_1$ 's head, and so on. If  $e_2$  is the edge that appears right below  $e_1$  on the path in  $F_{e_1}$ , then the path in  $F_{e_2}$  only touches the bottom two copies of  $G_{\ell,r}$  in the figure, and  $e_1$  is contained in a cycle in  $F_{e_2}$  (the one for the copy of  $G_{\ell,r}$  containing  $e_1$ ). This demonstrates the importance of including cycles in the frames.

We now proceed to enumerate all the cases to be checked. Let  $\ell$  be the level of  $e_1$ . If  $\ell < k$  then the path in  $F_{e_1}$  is comprised of seven segments: (i) edges at level  $\ell - 1$ , (ii) edges at level  $\ell$ , (iii) one edge at level  $\ell + 1$ , (iv) edges at level  $\ell$ , (v) one edge at level  $\ell + 1$ , (vi) edges at level  $\ell$ , and (vii) edges at level  $\ell - 1$ . If  $\ell = 1$  then segments (i) and (vii) are empty. If  $e_1$  is mediating then either segments (i) and (ii) are empty, or segments (vi) and (vii) are empty. If  $e_1$  is outer then segment (iv) is empty. The presence of empty segments does not change the argument, it just decreases the number of possibilities for  $e_2$ . The most general case, when all seven segments are nonempty, is illustrated in Figure 4. If  $\ell = k$  then the path

Figure 6: Frame for an edge at level  $\ell = k$ .

in  $F_{e_1}$  is comprised of three segments: (I) edges at level  $k - 1$ , (II) edges at level  $k$ , and (III) edges at level  $k - 1$ .

First suppose  $\ell < k$ . If  $e_2$  is in segment (i) then  $e_1$  is in  $e_2$ 's segment (iii). If  $e_2$  is in segment (ii) then  $e_1$  is in  $e_2$ 's segment (ii). If  $e_2$  is in segment (iii) then  $e_1$  is in  $e_2$ 's segment (i) (if  $\ell \leq k - 2$ ) or  $e_2$ 's segment (I) (if  $\ell = k - 1$ ). If  $e_2$  is in segment (iv) then  $e_1$  is in a cycle in  $F_{e_2}$ . If  $e_2$  is in segment (v) then  $e_1$  is in  $e_2$ 's segment (vii) (if  $\ell \leq k - 2$ ) or  $e_2$ 's segment (III) (if  $\ell = k - 1$ ). If  $e_2$  is in segment (vi) then  $e_1$  is in  $e_2$ 's segment (vi). If  $e_2$  is in segment (vii) then  $e_1$  is in  $e_2$ 's segment (v). Suppose  $e_2$  is in a cycle in  $F_{e_1}$ . If  $e_2$  is in a copy of  $G_{\ell,r}$  adjacent to the copy  $e_1$  is in, and  $e_2$  is pointing in the same direction as  $e_1$  (for example, the 2nd-from-top row of bold edges in Figure 4), then  $e_1$  is in  $e_2$ 's segment (iv). Otherwise,  $e_1$  is in a cycle in  $F_{e_2}$ .

Now suppose  $\ell = k$ . If  $e_2$  is in segment (I) then  $e_1$  is in  $e_2$ 's segment (iii). If  $e_2$  is in segment (II) then  $e_1$  is in  $e_2$ 's segment (II). If  $e_2$  is in segment (III) then  $e_1$  is in  $e_2$ 's segment (v). This finishes the proof of Claim 2.4, and hence of Lemma 2.2 and of Theorem 1.1.

### 3 Symmetric TSP Path Problem

In this section we prove Theorem 1.2. For every  $\varepsilon > 0$  we need to construct an instance  $d = (d_e)_{e \in E}$  on  $K_n = (V, E)$  with  $s, t \in V$  such that for some  $r = \gamma n$  with  $\gamma > 0$  depending on  $\varepsilon$ ,

$$\frac{\min_{x \in SP^{\text{int}}(K_n, s, t)} d \cdot x}{\min_{x \in N_+^r(SP(K_n, s, t))} d \cdot x} \geq 3/2 - \varepsilon. \quad (3.1)$$

For integers  $\ell \geq 1$  and  $q \geq 0$  we construct an undirected graph  $G_{\ell,q} = (V_{\ell,q}, E_{\ell,q})$  as follows (see the illustration in Figure 7). The graph  $G_{\ell,q}$  has two horizontal node-disjoint paths each with  $\ell$  edges. It also has two cliques each with  $3q + 3$  nodes, a left one and a right one. The cliques are node-disjoint from

each other and from the paths. The two left endpoints of the paths each have edges to all the nodes in the left clique, and the two right endpoints of the paths each have edges to all the nodes in the right clique. Finally, there is an additional node  $s$  with edges to all the nodes in the left clique, and an additional node  $t$  with edges to all the nodes in the right clique.

Also consider the graph  $G'_{\ell,q} = (V'_{\ell,q}, E'_{\ell,q})$  (illustrated in Figure 8) which is the same as  $G_{\ell,q}$  except that  $s$  and  $t$  are connected by a path with  $\ell$  new edges (using  $\ell - 1$  new nodes). We have  $V_{\ell,q} \subseteq V'_{\ell,q}$  and  $E_{\ell,q} \subseteq E'_{\ell,q}$ .

**Lemma 3.1.** *For all  $r \geq 0$ ,  $\ell \geq 1$ , and  $q \geq 0$  the following holds. For all  $x = (x_e)_{e \in E_{\ell,q}}$ , if  $x' = (x'_e)_{e \in E'_{\ell,q}}$  is the same as  $x$  but with all new edges  $e$  in  $E'_{\ell,q}$  having  $x'_e = 1$ , then  $x' \in N_+^r(ST(G'_{\ell,q}))$  implies  $x \in N_+^r(SP(G_{\ell,q}, s, t))$ .*

*Proof.* We prove the lemma by induction on  $r$ . Suppose  $r = 0$ . Trivially,  $x'(\delta_{G'_{\ell,q}}(v)) = 2$  for all  $v \in V_{\ell,q}$  implies  $x(\delta_{G_{\ell,q}}(v)) = 2$  for all  $v \in V_{\ell,q} \setminus \{s, t\}$  and  $x(\delta_{G_{\ell,q}}(v)) = 1$  for all  $v \in \{s, t\}$ . Now consider an arbitrary set  $\emptyset \subsetneq S \subsetneq V_{\ell,q}$  with  $|S \cap \{s, t\}| \neq 1$ , and assume without loss of generality that  $|S \cap \{s, t\}| = 0$ . Then  $x'(\delta_{G'_{\ell,q}}(S)) \geq 2$  implies  $x(\delta_{G_{\ell,q}}(S)) \geq 2$  since none of the new edges is in  $\delta_{G'_{\ell,q}}(S)$ . Now consider an arbitrary set  $\emptyset \subsetneq S \subsetneq V_{\ell,q}$  with  $|S \cap \{s, t\}| = 1$ . Then  $x'(\delta_{G'_{\ell,q}}(S)) \geq 2$  implies  $x(\delta_{G_{\ell,q}}(S)) \geq 1$  since only one new edge is in  $\delta_{G'_{\ell,q}}(S)$ . Thus we have shown that  $x' \in ST(G'_{\ell,q})$  implies  $x \in SP(G_{\ell,q}, s, t)$ .



Now suppose  $r > 0$  and the lemma holds for  $r - 1$ . Assume  $x' \in N_+^{r-1}(ST(G'_{\ell,q}))$  and thus it has a protection matrix  $X'$  (which has a 0th row and a 0th column, and the remaining rows and columns are indexed by  $E'_{\ell,q}$ ). Obtain a matrix  $X$  by deleting the rows and columns corresponding to the new edges in  $E'_{\ell,q}$ . We claim that  $X$  is a protection matrix witnessing that  $x \in N_+^{r-1}(SP(G_{\ell,q}, s, t))$ . Note that  $X$  is symmetric and positive semidefinite since it is a principal submatrix of the symmetric positive semidefinite matrix  $X'$ . We also have  $X_0 = \text{diag}(X) = (1 \ x)$ . We just need to verify that  $X_{e^*} \in \text{cone}(N_+^{r-1}(SP(G_{\ell,q}, s, t)))$  and  $X_0 - X_{e^*} \in \text{cone}(N_+^{r-1}(SP(G_{\ell,q}, s, t)))$  for each  $e^* \in E_{\ell,q}$ . We just consider  $X_{e^*}$ ; the case of  $X_0 - X_{e^*}$  is similar. We know that  $X'_{e^*} \in \text{cone}(N_+^{r-1}(ST(G'_{\ell,q})))$ . Thus if  $X'_{e^*,0} = 0$  then  $X'_{e^*}$  is all 0's, so  $X_{e^*}$  is all 0's and we have  $X_{e^*} \in \text{cone}(N_+^{r-1}(SP(G_{\ell,q}, s, t)))$ . Otherwise  $X'_{e^*,0} > 0$  and we have  $x'^{(e^*)} \in N_+^{r-1}(ST(G'_{\ell,q}))$  where  $x'^{(e^*)}$  is  $X'_{e^*}/X'_{e^*,0}$  with the 0th entry omitted. For each new edge  $e \in E'_{\ell,q}$ , since  $X'_{0,e} = x'_e = 1$  we must have  $X'_{e^*,e} = X'_{e^*,0}$ , by a basic property of protection matrices. Thus  $x'^{(e^*)}_e = 1$  for each new edge  $e$ , and the induction hypothesis tells us that  $x^{(e^*)} \in N_+^{r-1}(SP(G_{\ell,q}, s, t))$  where  $x^{(e^*)}$  is  $x'^{(e^*)}$  with the entries for the new edges omitted, or in other words,  $x^{(e^*)}$  is  $X_{e^*}/X_{e^*,0}$  with the 0th entry omitted. Thus we have  $X_{e^*} \in \text{cone}(N_+^{r-1}(SP(G_{\ell,q}, s, t)))$  as desired.  $\square$

Now fix some integers  $r \geq 0$  and  $\ell \geq 1$  and let  $q = r$  and  $n = |V_{\ell,r}| = 2(\ell + 1) + 2(3r + 3) + 2$ . We define the edge distances  $d = (d_e)_{e \in E}$  on the complete undirected graph  $K_n = (V_{\ell,r}, E)$  as the shortest path distances in the unweighted graph  $G_{\ell,r}$ . Note that  $d$  is nonnegative and satisfies the triangle inequality, so this is a valid instance (together with the distinguished nodes  $s, t$ ).

**Lemma 3.2.** *We have  $\min_{x \in SP^{int}(K_n, s, t)} d \cdot x \geq 3\ell - 2$ .*

**Lemma 3.3.** *We have  $\min_{x \in N_+^r(SP(K_n, s, t))} d \cdot x \leq 2\ell + 6r + 9$ .*

Combining Lemma 3.2 and Lemma 3.3, we find that the ratio on the left side of (3.1) is at least  $\frac{3\ell - 2}{2\ell + 6r + 9}$ , which is at least  $3/2 - \varepsilon$  provided  $\ell$  is large enough and  $r \leq \gamma'\ell$  for some small enough  $\gamma' > 0$

Figure 7: The graph  $G_{4,1}$ .Figure 8: The graph  $G'_{4,1}$ .

depending on  $\varepsilon$ . The latter is implied by  $r = \gamma n$  for some small enough  $\gamma > 0$  depending on  $\gamma'$ . It remains to prove Lemma 3.2 and Lemma 3.3.

*Proof of Lemma 3.2.* This is equivalent to lower-bounding the cost of a (non-simple)  $s$ - $t$  path in  $G_{\ell,r}$  that visits each node at least once. Given such a path, first note that at most one of the edges on the top and bottom paths in  $G_{\ell,r}$  is not traversed (otherwise some node would not be visited). We claim that for either the top path or the bottom path, all but at most one of the edges are traversed at least twice. Suppose not; then there exist two top edges and two bottom edges, all four of which are traversed at most once. By symmetry, assume the first of these four edges to be traversed is on top; then the  $s$ - $t$  path cannot return to the left clique without first going to the right clique. Once in the right clique, the path must traverse at least one of the two special bottom edges, but then there is no way to get back to the right clique to end at  $t$ . This proves the claim. These observations imply that the contribution of the top and bottom paths to the cost of the  $s$ - $t$  path is at least  $3\ell - 2$ .  $\square$

*Proof of Lemma 3.3.* Consider the vector  $x = (x_e)_{e \in E_{\ell,r}}$  defined as follows. The edges  $e$  on the two paths all have  $x_e = 1$ . The edges  $e$  within the two cliques all have  $x_e = \frac{2-1/(r+1)}{3r+2}$ . All remaining edges  $e$  have  $x_e = 1/(3r+3)$  (that is, those edges that go between a clique and an endpoint of one of the paths, or are incident to  $s$  or  $t$ ). Let  $x' = (x'_e)_{e \in E'_{\ell,r}}$  be the same as  $x$  but with all new edges  $e$  in  $E'_{\ell,r}$  having  $x'_e = 1$ . Cheung [11] proves that  $x' \in N_+^r(ST(G'_{\ell,r}))$ ; by Lemma 3.1 this implies  $x \in N_+^r(SP(G_{\ell,r}, s, t))$ . Then a simple argument shows that extending  $x$  with 0 values for all the edges in  $E \setminus E_{\ell,r}$  yields a point  $\hat{x} \in N_+^r(SP(K_n, s, t))$  such that  $d \cdot \hat{x} = \sum_{e \in E_{\ell,r}} x_e$  (using the fact that every edge in  $G_{\ell,r}$  is the shortest path

between its endpoints). We have

$$\sum_{e \in E_{\ell,r}} x_e = 2\ell \cdot 1 + 2 \binom{3r+3}{2} \cdot \frac{2-1/(r+1)}{3r+2} + 6(3r+3) \cdot \frac{1}{3r+3} = 2\ell + 6r + 9$$

so  $\hat{x}$  witnesses Lemma 3.3.  $\square$

## 4 Open Problems

There are many open problems on approximability and integrality gaps for variants of the traveling salesperson problem. There is much room for quantitative improvements in our results and those of [10]. Also, is the integrality gap lower bound of  $3/2$  (after one round) tight for the instance we use to prove Theorem 1.1? Can the lift-and-project procedures of Lovász-Schrijver or Sherali-Adams be employed to get improved algorithms for any variant of the traveling salesperson problem?

## Acknowledgments

I thank Luca Trevisan for suggesting to explore lift-and-project procedures applied to the traveling salesperson problem, and I thank Siu On Chan for discussions on lift-and-project procedures. I also thank anonymous reviewers for their comments.

## References

- [1] MIKHAIL ALEKHNOVICH, SANJEEV ARORA, AND IANNIS TOURLAKIS: Towards strong nonapproximability results in the Lovász-Schrijver hierarchy. *Computational Complexity*, 20(4):615–648, 2011. 3
- [2] HYUNG-CHAN AN, ROBERT KLEINBERG, AND DAVID SHMOYS: Improving Christofides’ algorithm for the  $s$ - $t$  path TSP. In *Proceedings of the 44th ACM Symposium on Theory of Computing*, pp. 875–886, 2012. 2
- [3] DAVID APPLEGATE, ROBERT BIXBY, VASEK CHVATAL, AND WILLIAM COOK: *The Traveling Salesman Problem: A Computational Study*. Princeton University Press, 2007. 2
- [4] SANJEEV ARORA: Polynomial time approximation schemes for Euclidean traveling salesman and other geometric problems. *Journal of the ACM*, 45(5):753–782, 1998. 2
- [5] SANJEEV ARORA, MICHELANGELO GRIGNI, DAVID KARGER, PHILIP KLEIN, AND ANDRZEJ WOŁOSZYN: A polynomial-time approximation scheme for weighted planar graph TSP. In *Proceedings of the 9th ACM-SIAM Symposium on Discrete Algorithms*, pp. 33–41, 1998. 2
- [6] ARASH ASADPOUR, MICHEL GOEMANS, ALEKSANDER MADRY, SHAYAN OVEIS GHARAN, AND AMIN SABERI: An  $O(\log n / \log \log n)$ -approximation algorithm for the asymmetric traveling salesman problem. In *Proceedings of the 21st ACM-SIAM Symposium on Discrete Algorithms*, pp. 379–389, 2010. 2, 3

- [7] SIAVOSH BENABBAS, KONSTANTINOS GEORGIOU, AVNER MAGEN, AND MADHUR TULSIANI: SDP gaps from pairwise independence. *Theory of Computing*, 8(1):269–289, 2012. 3
- [8] MOSES CHARIKAR, MICHEL GOEMANS, AND HOWARD KARLOFF: On the integrality ratio for the asymmetric traveling salesman problem. *Mathematics of Operations Research*, 31(2):245–252, 2006. 3, 8, 9, 10, 11
- [9] MOSES CHARIKAR, KONSTANTIN MAKARYCHEV, AND YURY MAKARYCHEV: Integrality gaps for Sherali-Adams relaxations. In *Proceedings of the 41st ACM Symposium on Theory of Computing*, pp. 283–292, 2009. 3
- [10] JOSEPH CHERIYAN, ZHIHAN GAO, KONSTANTINOS GEORGIOU, AND SAHIL SINGLA: On integrality ratios for asymmetric TSP in the Sherali-Adams hierarchy. In *Proceedings of the 40th International Colloquium on Automata, Languages and Programming*, pp. 340–351, 2013. 3, 9, 18
- [11] KEVIN CHEUNG: On Lovász-Schrijver lift-and-project procedures on the Dantzig-Fulkerson-Johnson relaxation of the TSP. *SIAM Journal on Optimization*, 16(2):380–399, 2005. 3, 4, 7, 9, 17
- [12] EDEN CHLAMTAC AND MADHUR TULSIANI: Convex relaxations and integrality gaps. In *Handbook on Semidefinite, Conic and Polynomial Optimization*. Springer, 2011. 3, 8
- [13] NICOS CHRISTOFIDES: Worst-case analysis of a new heuristic for the travelling salesman problem. Technical Report 388, Graduate School of Industrial Administration, Carnegie Mellon University, 1976. 2
- [14] GEORGE DANTZIG, DELBERT FULKERSON, AND SELMER JOHNSON: Solution of a large-scale traveling-salesman problem. *Journal of the Operations Research Society of America*, 2:393–410, 1954. 2, 5
- [15] URIEL FEIGE AND MOHIT SINGH: Improved approximation ratios for traveling salesperson tours and paths in directed graphs. In *Proceedings of the 10th International Workshop on Approximation Algorithms for Combinatorial Optimization Problems*, pp. 104–118, 2007. 2
- [16] ZACHARY FRIGGSTAD, ANUPAM GUPTA, AND MOHIT SINGH: An improved integrality gap for asymmetric TSP paths. In *Proceedings of the 16th Conference on Integer Programming and Combinatorial Optimization*, pp. 181–192, 2013. 3
- [17] ZACHARY FRIGGSTAD, MOHAMMAD SALAVATIPOUR, AND ZOYA SVITKINA: Asymmetric traveling salesman path and directed latency problems. *SIAM Journal on Computing*, 42(4):1596–1619, 2013. 3
- [18] ZHIHAN GAO: An LP-based  $3/2$ -approximation algorithm for the  $s$ - $t$  path graph traveling salesman problem. *Operations Research Letters*, 41(6):615–617, 2013. 2, 3
- [19] KONSTANTINOS GEORGIOU, AVNER MAGEN, TONIANN PITASSI, AND IANNIS TOURLAKIS: Integrality gaps of  $2 - o(1)$  for vertex cover SDPs in the Lovász-Schrijver hierarchy. *SIAM Journal on Computing*, 39(8):3553–3570, 2010. 3

- [20] ANUPAM GUPTA, KUNAL TALWAR, AND DAVID WITMER: Sparsest cut on bounded treewidth graphs: Algorithms and hardness results. In *Proceedings of the 45th ACM Symposium on Theory of Computing*, pp. 281–290, 2013. [3](#)
- [21] GREGORY GUTIN AND ABRAHAM PUNNEN, editors. *The Traveling Salesman Problem and Its Variations*. Springer, 2007. [2](#)
- [22] MICHAEL HELD AND RICHARD KARP: The traveling-salesman problem and minimum spanning trees. *Operations Research*, 18:1138–1162, 1970. [2](#), [5](#)
- [23] MICHAEL HELD AND RICHARD KARP: The traveling-salesman problem and minimum spanning trees: Part II. *Mathematical Programming*, 1:6–25, 1971. [2](#), [5](#)
- [24] HAN HOOGEVEEN: Analysis of Christofides' heuristic: Some paths are more difficult than cycles. *Operations Research Letters*, 10(5):291–295, 1991. [2](#)
- [25] MAREK KARPINSKI, MICHAEL LAMPIS, AND RICHARD SCHMIED: New inapproximability bounds for TSP. In *Proceedings of the 24th International Symposium on Algorithms and Computation*, pp. 568–578, 2013. [2](#)
- [26] PHILIP KLEIN: A linear-time approximation scheme for TSP in undirected planar graphs with edge-weights. *SIAM Journal on Computing*, 37(6):1926–1952, 2008. [2](#)
- [27] MICHAEL LAMPIS: Improved inapproximability for TSP. In *Proceedings of the 15th International Workshop on Approximation Algorithms for Combinatorial Optimization Problems*, pp. 243–253, 2012. [2](#)
- [28] EUGENE LAWLER, JAN LENSTRA, ALEXANDER RINNOOY KAN, AND DAVID SHMOYS, editors. *The Traveling Salesman Problem: A Guided Tour of Combinatorial Optimization*. Wiley, 1985. [2](#)
- [29] LÁSZLÓ LOVÁSZ AND ALEXANDER SCHRIJVER: Cones of matrices and set-functions and 0-1 optimization. *SIAM Journal on Optimization*, 1:166–190, 1991. [3](#), [4](#)
- [30] JOSEPH MITCHELL: Guillotine subdivisions approximate polygonal subdivisions: A simple polynomial-time approximation scheme for geometric TSP,  $k$ -MST, and related problems. *SIAM Journal on Computing*, 28(4):1298–1309, 1999. [2](#)
- [31] TOBIAS MÖMKE AND OLA SVENSSON: Approximating graphic TSP by matchings. In *Proceedings of the 52nd IEEE Symposium on Foundations of Computer Science*, pp. 560–569, 2011. [2](#)
- [32] MARCIN MUCHA: 13/9-approximation for graphic TSP. In *Proceedings of the 29th International Symposium on Theoretical Aspects of Computer Science*, pp. 30–41, 2012. [2](#)
- [33] VISWANATH NAGARAJAN: On the LP relaxation of the asymmetric traveling salesman path problem. *Theory of Computing*, 4(1):191–193, 2008. [7](#)

- [34] VISWANATH NAGARAJAN AND R. RAVI: The directed minimum latency problem. In *Proceedings of the 11th International Workshop on Approximation Algorithms for Combinatorial Optimization Problems*, pp. 193–206, 2008. [3](#)
- [35] THÀNH NGUYEN: A simple LP relaxation for the asymmetric traveling salesman problem. In *Proceedings of the 11th International Workshop on Approximation Algorithms for Combinatorial Optimization Problems*, pp. 207–218, 2008. [8](#)
- [36] SHAYAN OVEIS GHARAN AND AMIN SABERI: The asymmetric traveling salesman problem on graphs with bounded genus. In *Proceedings of the 22nd ACM-SIAM Symposium on Discrete Algorithms*, pp. 967–975, 2011. [2](#)
- [37] SHAYAN OVEIS GHARAN, AMIN SABERI, AND MOHIT SINGH: A randomized rounding approach to the traveling salesman problem. In *Proceedings of the 52nd IEEE Symposium on Foundations of Computer Science*, pp. 550–559, 2011. [2](#)
- [38] CHRISTOS PAPADIMITRIOU AND SANTOSH VEMPALA: On the approximability of the traveling salesman problem. *Combinatorica*, 26(1):101–120, 2006. [2](#)
- [39] GRANT SCHONEBECK, LUCA TREVISAN, AND MADHUR TULSIANI: Tight integrality gaps for Lovász-Schrijver LP relaxations of vertex cover and max cut. In *Proceedings of the 39th ACM Symposium on Theory of Computing*, pp. 302–310, 2007. [3](#)
- [40] ANDRÁS SEBÓ: Eight-fifth approximation for the path TSP. In *Proceedings of the 16th Conference on Integer Programming and Combinatorial Optimization*, pp. 362–374, 2013. [2, 3](#)
- [41] ANDRÁS SEBÓ AND JENS VYGEN: Shorter tours by nicer ears: 7/5-approximation for graphic TSP, 3/2 for the path version, and 4/3 for two-edge-connected subgraphs. *CoRR*, abs/1201.1870, 2012. [2, 3](#)
- [42] HANIF SHERALI AND WARREN ADAMS: A hierarchy of relaxations between the continuous and convex hull representations for zero-one programming problems. *SIAM Journal on Discrete Mathematics*, 3(3):411–430, 1990. [3, 5](#)
- [43] DAVID SHMOYS AND DAVID WILLIAMSON: Analyzing the Held-Karp TSP bound: A monotonicity property with application. *Information Processing Letters*, 35(6):281–285, 1990. [3](#)
- [44] MADHUR TULSIANI: Lovász-Schrijver reformulation. In *Wiley Encyclopedia of Operations Research and Management Science*. Wiley, 2011. [3](#)
- [45] LAURENCE WOLSEY: Heuristic analysis, linear programming and branch and bound. *Mathematical Programming Study*, 13:121–134, 1980. [3](#)

THOMAS WATSON

AUTHOR

Thomas Watson  
postdoctoral researcher  
University of Toronto, Toronto, ON  
thomasw@cs.toronto.edu  
<http://www.cs.toronto.edu/~thomasw>